

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

"Made available under NASA sponsorship
in the interest of early and wide dis-
semination of Earth Resources Survey
Program information and without liability
for any use made thereof."

7.6-10441
CR-148518

ESTUARINE DENSITY FRONTS AND
THEIR EFFECT ON OIL SLICKS

V. Klemas, D. Polis, G. Davis
College of Marine Studies *etc.*
University of Delaware

July 23, 1976
Report on Significant Results
NASA LANDSAT CONTRACT NAS5-20983
UN 20570

(E76-10441) ESTUARINE DENSITY FRONTS AND
THEIR EFFECT ON OIL SLICKS (Delaware Univ.)
2 p HC \$3.50 CSCL 13B

N76-29671

Unclas
G3/43 00441

Prepared for
GODDARD SPACE FLIGHT CENTER
Greenbelt, MD 20771

Significant Results

Estuarine fronts represent regions of extremely high gradient or discontinuity in various parameters of physical interest, the most important being the water velocity and density fields. Such fronts strongly influence pollutant dispersion, by capturing oil slicks and other pollutants concentrated in surface films and drawing them down into the water column. Aircraft and boats were combined to study the behavior of different types of fronts in Delaware Bay and their effect on pollutants in order to provide a basis for improving an oil drift and spreading model. Imagery from the LANDSAT satellites provided the most effective means of determining the location and extent of frontal systems over all portions of the tidal cycle. This information is being used to modify the oil drift and spreading model.